

CORRECTED  
VERSION\*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/10, C12Q 1/68		A3	(11) International Publication Number: WO 98/51789
			(43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/DK98/00186		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 13 May 1998 (13.05.98)		Published With international search report.	
(30) Priority Data: 0547/97 13 May 1997 (13.05.97) DK 08/871,030 19 May 1997 (19.05.97) US 0432/98 27 March 1998 (27.03.98) DK		(88) Date of publication of the international search report: 18 March 1999 (18.03.99)	
(71) Applicant (for all designated States except US): DISPLAY SYSTEMS BIOTECH APS [DK/DK]; Lersø Park Allé 40, DK-2100 Copenhagen Ø (DK).			
(72) Inventor; and (75) Inventor/Applicant (for US only): WARTHOF, Peter, Rolf [DK/DK]; Borgmester Jensens Allé 31B, DK-2100 Copenhagen Ø (DK).			
(74) Agent: PLOUGMANN, VINGTOFT & PARTNERS A/S; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).			

(54) Title: A METHOD TO CLONE mRNAs AND DISPLAY OF DIFFERENTIALLY EXPRESSED TRANSCRIPTS (DODET)

(57) Abstract

The present invention relates to a method for preparing a normalized sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample. It is an object of the present invention to provide new methods and means for investigating the expression patterns in cells, especially in eukaryotic cells. The results of such investigations may be used in drug development, gene discovery, diagnosis of diseases, etc.



\*(Referred to in PCT Gazette No. 49/1999, Section II)

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## A METHOD TO CLONE mRNAs AND DISPLAY OF DIFFERENTIALLY EXPRESSED TRANSCRIPTS (DODET)

## BACKGROUND OF THE INVENTION

The human body is comprised primarily of specialised cells performing different physiological functions organised into organs and tissues. All human cells contain DNA, arranged in a series of sub-units known as genes. It is estimated that there are approximately 100,000 genes in the human genome. Genes are the blueprints for proteins. Proteins may perform a wide variety of biological functions, for example messengers, catalysts and sensors. Such compounds are responsible for managing most of the physiological and biochemical functions in humans and all other living organisms. Over the last few decades, there has been a growing recognition that many major diseases have a genetic basis. It is now well established that genes play an important role in cancer, cardiovascular diseases, psychiatric disorders, obesity, and metabolic diseases. Significant resources are being focused on genomic research based on the notion that the nucleotide sequences of a particular gene and its predicted protein product will lead to an understanding of its function in healthy and malfunctioning cells or tissues. This understanding is expected, in turn, to lead to therapeutic and diagnostic approaches, focused on molecular targets associated with the gene and the protein it expresses. The first step on the way to the development of such applications is to identify the genes specifically involved in the different categories of diseases. Application of this knowledge can produce new and valuable markers, identifying regions producing major diseases to be used for diagnostic and therapeutic benefit.

Faced with the high complexity of the human genome, many approaches are being used to unravel the connection between primary gene structure and function. One well publicised approach is embodied in the Human Genome Mapping Project, where the sequence of all the individual genes in the entire human genome is painstakingly being determined. At the pre-

- sent, however, little information can be directly retrieved on the function of the identified genes and still less about temporal and spatial expression patterns of the developing or mature organism. Other approaches, such as random cDNA sequencing, involve the sequence determination of all genes expressed in a certain tissue, or developmental stage, of an organism. Like a number of other strategies, this is time consuming and prone to numerous problems.
- Although the flood of data from large scale sequencing programmes is of enormous benefit to the scientific community, one of the major problems faced by such "shotgun" approaches is the lack of specific information that can be retrieved without significantly more work on the biology of each of the individual genes.
- Several other approaches have been taken by molecular biologists to obtain more specific information on the genetic background of particular biological processes. Such approaches rely on a common concept. One gene, or a subset of genes, is switched on, initiating the healthy, pathological, or developmental status of an organ or cell type.

In a large number of experimental systems the isolation of genes, on the basis of their differential expression, has been applied successfully. Differential screening and subtractive hybridisation of cDNA libraries have become well established, cf. Zimmerman et al. (1980) and Davis et al. (1979). Differential library screening works well in practice for genes that are highly expressed, but mRNAs of low abundance are difficult to isolate. Subtractive hybridisation provides a more sensitive screening, but requires large amounts of RNA. More recently RNA fingerprinting methods (often referred to as differential display or DD/RT PCR) have been added to these tools, offering attractive new features for isolating genes. RNA fingerprinting methods are PCR based and therefore do not require large amounts of RNA for experiments. In addition to this, RNA fingerprinting methods allow a large number of RNA pools to be screened for specific

mRNAs simultaneously. Investigation of a wide range of pathogenic developmental stages and their controls would be possible. To date, two methods of RNA fingerprinting have proven useful for isolating genes. In 1992 Liang et al. published a  
5 protocol (US Patent 5,262,311), soon after a protocol from Welsh et al. (1992) was presented. Both methods begin with cDNA synthesis from RNA using at least one arbitrary primer for the initiation of first and second strand synthesis.

Welsh et al. (1992) designed a protocol in which the same  
10 arbitrary 20-mer oligo is used for first and second strand synthesis. Using arbitrary primers only a subset of the mRNAs are transcribed to cDNA. The cDNA pools are then used for a standard PCR with the same primers. One of the dNTPs in the PCR mix contains a radioactive label ( $^{35}\text{S}$  or  $^{32}\text{P}$ ) for visualisation of the PCR fragments with PAGE. The Liang and Welsh  
15 methods rely on at least one small arbitrary primer for selection of specific cDNAs. As a consequence annealing temperatures are low ( $\sim 40^\circ\text{C}$ ), and all amplified cDNA fragments originate from a certain degree of mismatch priming.  
20 Later several groups produced refinements and optimisations leading to a plethora of articles describing the usefulness of the method (Bauer and Warthoe et al. 1993; Warthoe et al. 1995; Liang and Warthoe et al. 1995; Rohde and Warthoe et al. 1996).


## 25 OBJECT OF THE INVENTION

It is an object of the present invention to provide new methods and means for investigating the expression patterns in cells, especially in eukaryotic cells. The results of such investigations may be used in drug development, gene discovery,  
30 ry, diagnosis of diseases etc., and therefore such improved methods are highly desirable.

## SUMMARY OF THE INVENTION


In its broadest scope, the invention pertains to a method for preparing a sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample, the  
5 method comprising the steps of

a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer having the general formula



10 wherein  $\text{Con}_1$  is any sequence between 1-100 nucleotides, dT is deoxythymidynyl, V is A, G or C, N is A, G, C or T,  $n2$  is an integer  $\geq 1$ ,  $n3$  is 0 or 1, if  $n3$  is 0 then  $n4$  is 0, and if  $n3$  is 1  $n4$  is an integer  $\geq 0$ , thereby obtaining first strand cDNA fragments,


15 b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, and a second cDNA primer with the general formula



20 wherein  $\text{Con}_2$  is any sequence between 1-100 nucleotides and can be different or identical to  $\text{con}_1$ ,  $\text{N}_x$  is A, G, T or C, and  $x$  is an integer  $\geq 0$ , in a appropriate enzyme/-buffer solution which comprises the DNA pol I enzyme or the Klenow fragment of the DNA pol I enzyme, all four  
25 deoxyribonucleoside triphosphates and standard buffer and temperature conditions, thereby obtaining double stranded cDNA fragments,

c) subjecting the cDNA fragments obtained in step b) to a molecular amplification procedure so as to obtain ampli-

fied cDNA fragments, wherein is used a set of amplification primers having the general formula



I

5 wherein  $\text{Con}_3$  is a sequence identical to either  $\text{Con}_1$  or  $\text{Con}_2$  or both, N is A, G, T or C, and  $n1$  is an integer  $\geq 0$ , wherein at least one set of primers has the general formula I where  $n > 0$ , said at least one set being capable of priming amplification of any nucleotide sequence complementary in its 5'-end to  $\text{Con}_1$  or  $\text{Con}_2$ .

10 This method is advantageous for amplifying very small amounts of RNA. Using the method of the invention it is possible to perform gene-profile analysis from less than 100 cells equal to  $10^{-9}$  gram total RNA (10 pgram RNA per cell).

15 In a further aspect, the invention relates to a method for preparing a sub-divided library of amplified cDNA fragments from the coding region of mRNA (which may be of prokaryotic, Archae or eukaryotic origin) contained in a sample, the method comprising the steps of

20 a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer, thereby obtaining first strand cDNA fragments,


25 b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, thereby obtaining double stranded cDNA fragments,

c) digesting the double stranded cDNA fragments with at least one restriction endonuclease, thereby obtaining cleaved cDNA fragments,



d) ligating at least two adapter fragments to the cleaved cDNA fragments obtained in step c), so as to obtain ligated cDNA fragments, and

5 e) subjecting the ligated cDNA fragments obtained in step d) to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used, for an adapter fragment used in step d), a set of amplification primers having the general formula



II

10 wherein Com is a sequence complementary to at least the 5'-end of an adapter fragment which is ligated to the 3'-end of a cleaved cDNA fragment, N is A, G, T, or C, and  $n1$  is an integer  $\geq 0$ , and wherein at least one set of primers has the general formula II where  $n1 > 0$ , said at  
15 least one set being capable of priming amplification of any nucleotide sequence ligated in its 3'-end to the adapter fragment complementary in its 5'-end to Com.

The overall advantage of the invention compared to the prior art is that the resulting library of cDNA fragments contains  
20 nucleic acid sequences from all parts of cDNA which is produced in step a). Prior art techniques which *i.a.* rely on poly-dT cDNA priming have a tendency to only yield fragments derived from the long untranslated regions of mRNA. Furthermore, by fine-tuning of the conditions in each step, the  
25 method of the present invention results in highly specific reproduction of sequence information which is present in mRNA, even in mRNA which is only present in relatively low amounts. Furthermore, by choosing the optimum composition of endonuclease(s) it is possible to obtain cDNA fragments which  
30 are derived from a very large percentage of the total number of transcribed genes in relevant cells.

The present method allows the targeted visualisation of known genes by using primer combinations, corresponding to sequen-



ces from the gene of interest. This has the advantage that all steps of the procedure and the biological system can easily be verified. Also, very specific expression analyses can be carried out on related genes with very high homology which could not be achieved by using hybridisation technology.

Briefly, further steps in the method of the invention involve isolation of bands of interest from a gel, their cloning and sequencing. The sequence information allows re-amplification of individual bands, using primers with the appropriate 3-4 nucleotide extensions. When run on a gel, these reactions will show one, or only a few, bands per lane, giving an unequivocal determination of band identity.

Since the present technology makes use of end labelled primers for visualization, the technology can be used, both with standard technologies involving radioactivity, or with fluorescent labelled primers, without the need for further optimisation.

The invention also pertains to methods for detecting differences between expression level(s) in cells which have been subjected to different conditions, methods for diagnosing disease, and methods related to "bioinformatics" wherein are used a combination of output from the above-disclosed method and data obtained by computer-simulation of corresponding treatment of well-defined stretches of nucleic acids.

A separate part of the invention pertains to a novel method for performing reverse transcription, methods which yield considerably enhanced quality in the reversely transcribed material. Also means for carrying out this separate part of the invention are disclosed.

In the following is given a short discussion of terms used in the present application:

"A sub-divided library of amplified cDNA fragments" is in the present context a library of amplified cDNA fragments which is split into a number of separate pools, each pool being defined by the sequences of the termini of the amplified  
5 fragments. For example, one pool may contain amplified fragments which are all characterized by having the sequence 5'-Com-AGC- in one of the strands, whereas another pool contains amplified fragments having the sequence 5'-Com-AAT in one of the strands. For a discussion of the meaning of "Con" and  
10 "Com", cf. below.

"A normalised library" is a library containing substantially equal representation of each mRNA, i.e. approximately the same number of copies of each mRNA.

"Reverse transcription" has its usual meaning in the art,  
15 i.e. synthesis of DNA using RNA as a template and effected by an enzyme having reverse transcriptase activity.

"Adapter fragment" is intended to mean a nucleic acid sequence containing a known sequence which can be used as template for a primer in a subsequent molecular amplification procedure such as PCR. The adapter fragment is further characterized by its ability to become integrated at the end of a  
20 cDNA fragment which has previously been cleaved with a restriction endonuclease in step c). In most cases, the restriction endonuclease leaves fragments having "sticky ends",  
25 to which the adapter fragment will anneal readily, and thereafter the adapter fragment becomes ligated to the cDNA by the action of a DNA ligase.

#### DETAILED DISCLOSURE OF THE INVENTION

In the following, the impact of each of the steps will be  
30 discussed in detail, see Figures 1 and 7.

Step a)

The goal in step a) is to produce a mixture of first strand cDNA fragments which is optimized in its composition for carrying out the subsequent steps. A number of considerations  
5 apply:

First of all, to reduce the "background noise", it is preferred that the annealing of cDNA primer to the RNA in step a) is performed under high stringency conditions, thereby ensuring that a minimum of mismatches are introduced in the  
10 cDNA relative to the mRNA, i.e. at a temperature above 50°C.

Secondly, it is desirable to obtain copies of sequences which are derived from all parts of mRNA in order to obtain information relating to the translated part of the mRNA. Prior art methods for reverse transcription of eukaryotic material have  
15 often utilised poly-dT as cDNA primers. This strategy has, however, the disadvantage that the most efficiently reverse transcribed material is situated in the untranslated part of the genes of interest. Hence, the only parts of the mRNA which become "visible" after e.g. a PCR procedure will very  
20 often be derived from untranslated regions of the RNA. The reason for this is two effects. First of all, the poly-dT approach has the consequence that the initiation point of reverse transcription is situated very far from e.g. the start codon relating to the operon in question. Secondly, the  
25 mRNA may include structures (e.g. "hairpin" structures due to intra-chain base-pairing) which block reverse transcription and by always initiating reverse transcription at one terminus of a gene, such structures will statistically block reverse transcription of a number of translated regions.

30 It is in the present invention preferred to ensure that cDNAs are produced in step a) which are representatives of the entire gene, including the translated regions.

This can be obtained in a number of ways. If poly-dT priming (or a variation thereof) is used, it is preferred to perform the reverse transcription at an elevated temperature, e.g. in the range from about 45°C to about 95°C, and to use an enzyme  
5 having reverse transcriptase activity at said temperature. Normally the temperature will be higher than 45°C, e.g. at least 50°C, or even higher, e.g. at least 55°C, at least 60°C, at least 65°C or even higher, e.g. at least 70°C. This approach has the effect that the elevated temperature ensures  
10 that e.g. hairpin structures are "stretched out" during the reverse transcription step, thereby avoiding the lack of reversely transcribed fragments upstream of such structures.

Known enzymes having reverse transcriptase activity at such elevated temperatures are enzymes selected from the group  
15 consisting of DNA polymerases derived from thermophilic eubacteria, such as the polymerases Taq (Thermus aquaticus), Stoffel (Thermus aquaticus), Tht (Thermus thermophilus), Tfl/Tub (Thermus flavus), Tru (Thermus Ruber), Tca (Thermus caldophilus), Tfil (Thermus filiformis), Tbr (Thermus Brocki-  
20 anus), Bst (B. Stearothermophilus), Bca (B. Caldopenax YT-G), Bcav (B. Caldovelox YT-F), FjSS3-B.1 (Thermotoga FjSS3-B.1), Tma (Thermus Maritima), UITma (T. Maritima), Tli (T. Litoralis), Tli exo- (T. Litoralis), 9°N-7 (Thermococcus sp.), BG-D (Pyrococcus sp.), Pfu (P. furiosus), Pwo (P. woesei), Sac (S.  
25 Acidocaldarius), SsoI (S. Solfataricus), Tac (T. Acidophilum), and Mth (Methanococcus Voltae).

One minor disadvantage of using these thermostable enzymes is that they have a tendency to be relatively ineffective compared to the "traditional" non-thermostable, reverse tran-  
30 scriptases. Hence, especially if priming of the reverse transcription is not limited to the use of poly-dT primers, it is according to the invention possible to use non-thermostable, reverse transcriptases. Hence, in other preferred embodiments, the reverse transcription is carried out at a  
35 temperature in the range from about 25°C to about 55°C by use of an enzyme having reverse transcriptase activity at said

temperature. Normally the temperature will not exceed 50°C, and usually it will be lower, such as at most 47°C, at most 45°C, at most 43°C, at most 40°C, and at most 35°C. The reverse transcriptase can e.g. be selected from the group  
5 consisting of the reverse transcriptases from AMV (Avian Myeloblastosis Virus), M-MuLV (murine M-MuLV pol gene), and HIV-1 (HIV virus).

According to the invention, the most preferred way of carrying out step a) is to carry out reverse transcription in two  
10 subsequent steps, the first step comprising carrying out reverse transcription at the temperature conditions described above for non-thermostable enzymes, and the second step comprising carrying out reverse transcription at the temperature conditions described above for thermostable enzymes.  
15 Normally this can be accomplished by having two non-identical enzymes present in the reverse transcription reaction, especially because the non-thermostable enzyme will be inactivated by the increase in temperature which is introduced when going into step 2. Of course, the enzymes can be added for  
20 each reaction step, but it is preferred that both enzymes are present from the start of the reaction.

It is especially preferred that the activity of the enzyme which is active in the first step is substantially abolished in the second step (e.g. as a consequence of temperature  
25 denaturing of that enzyme), or expressed otherwise, that in the second step the enzyme used in the first step is substantially inactive. In general, it is preferred that the enzymes used in each step are substantially more active in the relevant temperature range than the one wherein the other enzyme  
30 is used.

In a preferred embodiment the reaction mixture with the sample comprises a cDNA primer, said cDNA primer being sufficiently complementary to the target RNA present in the sample to hybridize therewith and initiate synthesis of a single  
35 stranded cDNA molecule complementary to said target RNA and

the reaction mixture comprises an appropriate buffer which comprises all four deoxyribonucleoside triphosphates and a divalent cation selected from the group of  $Mg^{+2}$  and  $Mn^{2+}$  in a concentration between 0.1 and 5 mM.


- 5 In fact, it is believed that the above strategy for conducting reverse transcription by use of two enzymes having different temperature optima and of which one has a temperature optimum at which impeding structures in the RNA are "stretched out", is novel and inventive in its own right.
- 10 Preferred combinations of enzymes in this embodiment of the invention are that the enzyme effecting reverse transcription in the first step is MMuLV, AMV, HIV-1 and/or the enzyme effecting reverse transcription in the second step is Tth or Taq.
- 15 An object of the method of the invention is to obtain a subdivision of the cDNA produced. When the mRNA is derived from a eukaryotic system, the at least one cDNA primer may include an oligo or poly dT tail in the 5'-end, having the general formula  $5'-dT_{n2}-V_{n3}-N_{n4}-3'$ , wherein dT is deoxythymi-
- 20 danyl, V is A, G, or C, N is A, G, C, or T,  $n2$  is an integer  $\geq 1$ ,  $n3$  is 0 or 1, if  $n3$  is 0 then  $n4$  is 0, and if  $n3$  is 1 then  $n4$  is an integer  $\geq 0$ . It will be clear that when  $n3$  and  $n4$  are both zero, then the primer is an ordinary poly- or oligo-dT cDNA primer. However, when  $n3$  is 1, then the primer
- 25 is in fact a primer composition which will be able to prime the reverse transcription of any mRNA having a poly-A tail. If the original sample of RNA is subdivided, and each subpool is subjected to reverse transcription which uses one of the possible primers having the above formula where  $n3$  is 1,
- 30 then the result is a number of single stranded cDNA pools which are each different from each other in the 5'-end.

For example, when  $n3$  is 1,  $3 \times 4^{n4}$  groups of cDNA primers are used, each group being distinct from any one of the other groups with respect to the structure  $-V_{n3}-N_{n4}-$ . In such an



embodiment the pool of mRNA is conveniently subdivided into  $3 \times 4^{n4}$  aliquots which are each subjected separately to step a) utilising one of the  $3 \times 4^{n4}$  groups of cDNA primers, thereby obtaining a subdivision of the first strand cDNA into  $3 \times 4^{n4}$  separate pools. Normally  $n4$  will be 0 or 1, resulting in the provision of 3 or 12 pools, respectively.

When the starting material is not eukaryotic or when it is not the intention to necessarily set out from the part of the transcribed gene which is most remote relative to the translation start codon, the at least one cDNA primer does not include a poly or oligo dT tail in the 5'-end, or, alternatively, at least two cDNA primers are used of which at least one includes a poly or oligo dT tail in the 5'-end and of which at least one second does not include a poly or oligo dT tail in the 5'-end. Preferably, the cDNA primer which does not include a poly or oligo dT tail in the 5' end has the following structure



wherein N is A, G, T, or C, and x is an integer  $1 \leq x \leq 20$ . It will be clear that this corresponds to cDNA priming setting out from any translation stop codon. As for the above embodiments utilising a poly- or oligo-dT tailed primer, it is, by preparing primers having all possible permutations represented in the group  $N_x$ , possible to compose the primers so as to correspond to any possible sequence preceding a stop codon, thereby ensuring priming of all sequences having a stop codon in their sequence.

#### Step b)

This step is carried out by methods well known in the art. It is, however, preferred that step b) is carried out under conditions which minimize the formation of mismatches between nucleotides in the first and second cDNA strands. The double stranded cDNA procedure can be performed according to stan-



dard methods as described in Sambrook et al. (1989). However since standard polymerases can have difficulty in synthesizing regions containing secondary structures or with high GC-content, thermostable RNase H (Hybridase Thermostable RNase H, US 5,268,289) and thermostable rBst DNA polymerase from *Bacillus stearothermophilus* help overcome some of the limitations that standard polymerases (low temperature polymerases) suffer from.

Step c)

10 In one embodiment of the invention the ligated cDNA fragments obtained in step b) are subjected to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used a set of amplification primers having the general formula

15 
$$5' - \text{Con}_3 - \text{N}_{n1} - 3'$$
 I

wherein  $\text{Con}_3$  is a sequence identical to either  $\text{Con}_1$  or  $\text{Con}_2$  or both, N is A, G, T or C, and  $n1$  is an integer  $\geq 0$ , wherein at least one set of primers has the general formula I where  $n > 0$ , said at least one set being capable of priming amplification of any nucleotide sequence complementary in its 5'-end to  $\text{Con}_1$  or  $\text{Con}_2$ .

20

In another embodiment, after the preparation and optional subdivision of the mRNA, each of the different pools of cDNA is digested with at least one restriction enzyme to produce fragments of a size which can be separated using an appropriate size fractionation method.

25

The choice of restriction enzyme is based largely on the frequency of the cleavage sites in a given cDNA pool. Too many cleavage sites in each cDNA fragment will result in too small fragments, and vice versa. Optimally, the at least one enzyme should cleave every cDNA to yield fragments of the desired size. Statistically, it is not possible to cleave

30

every cDNA, but on the other hand a very large percentage can be cleaved by choosing a suitable enzyme or combination of enzymes. It is preferred that the method of the invention utilises at least one restriction enzyme chosen so as to  
5 ensure that at least 60% of cDNAs are cleaved, but higher percentages such as at least 65%, at least 70%, at least 75%, at least 80%, or even at least 85% are more preferred.

Preferably the invention should use restriction enzymes that leave protruding ends (sticky ends) at the termini of the DNA  
10 after digestion in step c), since this greatly facilitates the introduction of the adapter fragments in step d).

As will appear from the above, the frequency with which the restriction endonuclease cleaves is important. The at least one restriction enzyme is preferably chosen so as to cleave  
15 each complete cDNA into an average of about 3 fragments. It will be understood that some cDNAs obtained from preceding steps will not be cut at all (although this is a rare incidence when the restriction enzyme(s) is/are carefully chosen) whereas others will be cut with a high frequency. It has come  
20 out that use of a rare 4 base cutter as at the least one restriction endonuclease (such as the 4 base cutter AclI, AluI, BfaI, BstUI, Csp6I, DpnI, DpnII, HaeIII, HhaI, HinPII, HpaII, MboI, MnlI, MseI, MspI, NlaIII, RsaI, Sau3AI, TaiI, TaqI, and Tsp509I) ensures the optimum performance of the  
25 inventive method. By use of such a rare 4 base cutter, the use of only 1 restriction enzyme in step c) is sufficient and results in superior output.

Alternatively, a combination of restriction endonucleases can be used wherein a balance of e.g. 6 base cutters and 4 base  
30 cutters ensures a reasonable distribution of fragment sizes. For instance the use of a first restriction enzyme (e.g. a 6 base cutter) which statistically cleaves at least 20% of complete cDNA derived from the mRNA sample into two subfragments, and of a second restriction enzyme (e.g. a 4 base  
35 cutter) which statistically cleaves at least 50% of said

subfragments into 3 further subfragments, will also result in a series of fragments suitable for later size fractionation.

Step d)

The mixture(s) obtained in step c) are then subjected to a  
5 reaction wherein adapter fragments are added to both ends of the double stranded cDNA fragments obtained. As mentioned above, this part of the procedure is greatly facilitated by the cleaved cDNA fragments having protruding "sticky" ends, because pre-designed adapter fragments which fit to these  
10 protruding ends can easily be prepared.

The adapter (or anchor) fragments are added to the cleaved fragments in order to obtain "order in chaos" in the subsequent step. By adding known sequences to the termini of the cleaved fragments, one creates targets for specific amplifi-  
15 cation primers which can be designed specifically with the aim of amplifying sequences complying to the adapter fragments. The material thus obtained (primary template) can be pre-amplified, using primers complementary to the ligated adaptor sequences, giving rise to secondary template. The  
20 pre-amplification of primary template allows virtually unlimited amounts of template to be produced from one RNA preparation, avoiding the need for repeated isolations.

The adaptor sequence is thus selected so as to serve as the starting point for DNA polymerisation in e.g. a PCR reaction.  
25 The adaptor sequences are constructed in such a way that the specific endonuclease sites are not regenerated after ligation of said adaptor.

In a preferred embodiment at least one termination fragment is also ligated to the 3'-end of single strands of cleaved  
30 cDNA fragments, said at least one termination fragment introducing a block against DNA polymerization in the 5'→3' direction setting out from the at least one termination fragment and said at least one termination fragment being unable to

anneal to any primer of the at least two primer sets in step e) during the molecular amplification procedure.

The above is a very important procedure when combined with the use of detection effected by labelled primers in the amplification step, wherein only one member of the pair of primers is labelled whereas the other is designed to split up the amplified products according to their base composition adjacent to the adapter fragment. One important feature is that a single stranded cDNA fragment which has been provided with a termination fragment will not be amplified, because no primers will be able to anneal to the products of a first round polymerisation wherein such a fragment was the template, see Figure 7. Secondly, the approach opens for the possibility of removing background "noise" in a subsequent detection phase.

Normally, the at least one termination fragment comprises or is a chemically modified nucleotide sequence, such as for instance a nucleotide sequence which comprises a dideoxynucleotide in the 3'-end; this termination technique is well-known from e.g. the chain-termination sequencing technique according to Sanger. Under normal circumstances, the dideoxynucleotide should, according to the invention, be covalently attached to the nucleotide strand so as to avoid loss of the dideoxynucleotide during subsequent rounds of amplification. Superior stabilisation is attained if the dideoxynucleotide is phosphorylated.

As mentioned above, the ligation of adapter and/or termination fragments to the cleaved cDNA fragments in step d) is conveniently achieved by annealing the adapter fragments to sticky ends of the cDNA resulting from the cleavage in step c) and subjecting the product to the action of an enzyme having DNA ligase activity. Any suitable DNA ligase known in the art can be used.

Step e)

Step e) of the method of the invention results in the final sorting of the modified cDNA fragments from step d). As steps b), c) and d) are combined in the broadest embodiment of the invention, step e) corresponds to step c) of this embodiment described above.

The primers having the structure of formula I (step c) or II (step e) are designed so as to selectively amplify synthesized double stranded cDNA fragments obtained in step b) or predefined subsets of the adapted fragments obtained in step d). A number of ways this can be done may be envisaged, but the main strategy is to prime amplification in a series of separate reactions where the nucleotide sequence of one primer in one reaction ensures that the amplified products of that reaction are different from those obtained from any of the other reactions and that all the reactions result in amplification of all fragments obtained from step b) or d), respectively.

Even though the at least one set of amplification primers of formula I or II wherein has a  $n_1$  which is  $\geq 0$ , it is preferred that  $n_1=1$ ,  $n_1=2$ ,  $n_1=3$ , or  $n_1=4$  in one of the primers, because the number of primer fragments to be used in the reactions in order to cover all possible nucleotide stretches adjacent to the Con or adapter fragment is easily manageable. For instance, if  $n_1=5$ , it would be necessary to use  $4^5=1024$  different primers in order to obtain amplification of all possible nucleotide sequences adjacent to the relevant adapter fragment, and since the preferred embodiment of the invention requires that each such primer is used in a separate reaction, the work involved would be problematic.

It is also preferred that in one of the primers  $n=0$ , and it is especially preferred that this primer is labelled, in order to facilitate determination of the amplified fragments.

Hence, in the most preferred embodiments of the invention, the adapted cDNA fragments are amplified in a number of separate reactions wherein a labelled primer is used (which is normally identical in all reactions) and at least one non-labelled primer which is a member of the set of primers described above where  $n > 1$ . It is preferred that this set of amplification primers of formula I or II wherein  $n_1 > 0$  comprises all possible combinations and permutations of A, G, T, and C in the group  $N_{n_1}$ , since this will ensure that all possible cDNA fragments can be amplified by the set.

Hence, the ligated cDNA fragments are sub-divided into a number of pools prior to the molecular amplification in step e), and each pool is subjected to the amplification using a subset of the set of amplification primers, and in the most preferred embodiment the ligated cDNA fragments of step d) are subdivided into  $4^{n_1}$  pools which are each subjected separately to step e) wherein is used one amplification labelled primer as described above ( $n_1 = 0$ ) and one primer from the set of amplification primers as defined above ( $n > 0$ ), said one primer being distinct from any one of the primers used for amplifying any of the other pools. By using this approach, the originally reverse transcribed and cleaved cDNA fragments are subdivided into  $4^{n_1}$  pools which can each be subjected to further steps.

#### 25 Further steps and applications

The material obtained from the above-described series of reactions can now be utilised in a number of ways. Normally, a further step of separating amplified fragments obtained from the molecular amplification procedure is performed. This yields a mixture of amplified fragments which are separated e.g. by size separation, by mobility in a gel electrophoresis or by any suitable chromatographic method. Furthermore, a step of identification (e.g. by visualization of these separated fragments) is normally carried out for "book-keeping purposes"; the separated mixture of fragments will normally



be compared to some kind of reference which may be material derived from the same or another cell type.

Visualization of the separated fragments can, as mentioned above, be achieved by one of the primers in the amplification reaction being labelled, but other methods are of course available. For instance, a specifically labelled probe which e.g. binds to one of the adapter sequences will visualise the fragments, but also labelled nucleotides which have been incorporated in the fragments during the amplification procedure (e.g. a PCR) will of course be a suitable means for detection (e.g. by incorporating radioactive or fluorescent alpha dNTP into the cDNA fragment during PCR, where N = A, C, T, U or G).

However, it is preferred that visualisation of specific RNA Derived Fragments (RDFs) is achieved using primers which are radioactively or fluorescently labelled and are homologous to the adaptors. The comparatively high annealing temperatures (touch-down from 65°C to 56°C) which are preferably used ensure that polymerisation events will predominantly originate from perfect priming of adapter sequences and adjacent selective bases. Band intensities are largely a function of initial template concentration, whereas band intensities of the original Differential Display methods are dependent on the quality of the match between the individual template and primer. The visualisation of rare mRNAs using the present inventive methods will be less hampered by the over-representation of signal from highly abundant mRNAs. As in the case of arbitrary priming, the mismatch amplification and abundant RDFs always out-compete the amplification of rare fragments base pair perfectly. Our experiments suggest that as few as 100 molecules can be routinely detected in a given template. This corresponds to less than 1 transcript per cell in the original tissue.

One interesting part of the invention relates to the use of the above-described methods in bioinformatics. In short,

**RECTIFIED SHEET (RULE 91)**  
**ISA/EP**



known DNA sequences are inputted into a computer database, and on the basis of such sequences a comparison with a real-life run of the above-described methods can be performed. In this way, bands in a gel obtained from the methods of the invention can be unambiguously identified with respect to sequence, origin and even functionality. Hence, this part of the invention pertains to a method for determining the presence of an expression product in a cell or group of cells, the method comprising providing an RNA-containing sample from the cell or group of cells and subjecting the sample to the method described above, and thereafter performing a comparison of the thus identified amplified cDNA fragments with a database output, said database output comprising a computer-generated list of molecular weights of restriction DNA fragments of known sequences, said list being prepared by

- inputting and storing DNA sequence data in a database as virtual DNA sequences (these can be obtained and updated regularly from any database containing information about gene sequences from the relevant organism or cell type),
- 20 - subsequently simulating cleavage of the virtual DNA sequences with the at least one restriction nuclease and storing the resulting simulated cleavage products as virtual cleaved DNA fragments (such simulation is relatively uncomplicated, since the recognition and cleavage patterns of a large number of restriction enzymes are already known),
- 25 - simulating ligation to the virtually cleaved fragments of the at least two adapter fragments and storing the results as virtually ligated DNA fragments (again, this merely requires that input is provided of the structure of adapter fragments used in the real-life process),
- 30 - for each individual combination of primers used in step e) grouping the virtually ligated DNA fragments susceptible to amplification by said combination of primers in the same group,
- 35

- determining, in each group, the absolute and/or relative molecular weight of each virtually ligated DNA fragment, and
- outputting the content of each group in the form of a list comprising the absolute and/or relative molecular weights of the virtually ligated fragments in the group.

It is preferred that a link is maintained between each member of the output list and the original sequence from which such a member has been derived. This can e.g. be done by linking the input DNA sequence data to data relating to the genetic origin of the DNA sequence data and optionally to data relating to functional features relating to the genetic origin and thereafter maintaining the information as a pointer back in the system to said sequence. Hence, the output indication will conveniently further comprise information about the genetic origin of the virtually ligated DNA fragment and optionally information about functional features associated with the genetic origin.

For ease of use of such a bio-informatic system, it is normally necessary that 1) either the comparison is performed by inputting the identified amplified cDNA fragments in a format which allows automated comparison with the database output, or 2) the database output is outputted in a format which allows for direct comparison between the separated amplified cDNA fragments and the database output. For instance, if the visualized and separated cDNA fragments from step e) have been run on a gel, it will be possible to either read a digital reproduction of the gel pattern into the computer and let the computer compare this input with the computer generated pattern, or alternatively, to output the computer generated pattern in such a manner that it resembles an electrophoresis gel pattern.

Another part of the invention pertains to the use of the inventive method for comparing expression levels in different cells. One way of doing this is to determine the change in

expression, compared to the expression in a reference cell or reference group of cells, of an expression product in a cell or group of cells which has been subjected to a first set of conditions influencing the expression pattern of said cell or group of cells, said reference cell or group of cells being  
5 subjected to a second set of conditions, the method comprising providing an RNA-containing sample from the cell or group of cells and subjecting the sample to the method of the invention for sub-division, thereby obtaining data describing  
10 the amplified cDNA fragments derived from the sample, providing reference data describing amplified cDNA fragments derived from an RNA-containing reference sample from the reference cell or reference group of cells, the reference data being obtained by having previously subjected the reference  
15 sample to the method of the invention, subsequently performing a comparison of the data and the reference data to identify the cDNA fragments which are expressed at different levels in the two data sets, and thereafter using the differentially expressed cDNA fragments to determine which expression products are subject to a change in expression level. In  
20 other words, the method of the invention is carried out twice on the basis of two different RNA samples derived from cells subjected to differing conditions.

Normally, the data and reference data are selected from the  
25 group consisting of the apparent molecular weights of the amplified DNA fragments, the  $M_r$  of the amplified DNA fragments, the absolute amount of the amplified DNA fragments, and the relative amounts of the amplified DNA fragments. The reference data can further be extracted from a database  
30 containing the reference data defined above and optionally further information relating to the genetic origin of each amplified cDNA fragment from the reference.

Related to the above, the invention also allows for diagnosis  
of disease which is characterized by a deviating (increased  
35 or reduced) expression level of at least one expression product in at least one cell type, the method comprising pro-

viding an RNA-containing sample derived from the at least one cell type, subjecting the sample to the method of the invention thereby obtaining data describing the amplified cDNA fragments derived from the sample, providing reference data describing amplified cDNA fragments derived from a RNA-containing reference sample derived from the same type of cell from a subject not suffering from the disease, the reference data being obtained by having previously subjected the reference sample to the method according to the invention, and subsequently performing a comparison of the data and the reference data with respect to those cDNA fragments which are known to be related to the disease, and assessing whether a significant difference in the data and reference data exists so as to establish whether the expression level of the expression product deviates or not.

As for the embodiment above, also here the data and reference data are selected from the group consisting of the apparent molecular weights of the amplified DNA fragments, the  $M_r$  of the amplified DNA fragments, the absolute amount of the amplified DNA fragments, and the relative amounts of the amplified DNA fragments, and also here the reference data can be extracted from a database containing the reference data defined above and optionally further information relating to the genetic origin of each amplified cDNA fragment from the reference.

Further, the invention provides a method for treatment of a disease which is characterized by a deviating (increased or reduced) expression level of at least one expression product in at least one cell type, the method comprising providing an RNA-containing sample derived from the at least one cell type, subjecting the sample to the method of the invention thereby obtaining data describing the amplified cDNA fragments derived from the sample, providing reference data describing amplified cDNA fragments derived from a RNA-containing reference sample derived from the same type of cell from a subject not suffering from the disease, the reference

data being obtained by having previously subjected the reference sample to the method according to the invention, and subsequently performing a comparison of the data and the reference data with respect to those cDNA fragments which are  
5 known to be or suspected of being related to the disease, and assessing whether a significant difference in the data and reference data exists so as to establish whether the expression level of the expression product deviates or not.

If the expression product is reduced, the disease may be  
10 treated by delivering the expression product; if the expression product is increased, the disease may be treated by delivering an inhibitor (e.g. an antibody) against the expression product. The scope of the present invention includes an expression product identified by the method of the invention as such as well as methods for treating a disease which  
15 method has been provided by means of the method of the invention.

The mixtures of amplified fragments obtained from step e) of the method of the invention may also be used for preparing a  
20 surface (chip) coated with cDNA fragments. This can be done by

- subjecting an RNA-containing sample to the subdivision method of the invention including separation steps, and
- transferring the separated amplified cDNA fragments to a  
25 chip surface adapted to stably bind the separated amplified cDNA fragments while maintaining the spatial relative distribution pattern thereof.

Alternatively, such a chip can be prepared by

- subjecting an RNA-containing sample to the method of the  
30 invention without performing the separation, and thereafter

separating, by electrophoresis, the thus obtained amplified cDNA fragments on a particular surface adapted to stably bind the separated amplified cDNA fragments while maintaining the relative distribution pattern after electrophoresis. In this embodiment, the electrophoresis is preferably in the form of microelectrophoresis.

Transfer to the surface is preferably accomplished by a electrophoretic blotting technique, and/or by well-known photo-activated organic or inorganic chemistry coupling techniques.

The invention also pertains to a surface obtainable by the above-mentioned method for the preparation thereof. Such surfaces are considered novel and inventive, since known "DNA chips" rely on specific introduction of an array of nucleic acid fragments of known structure, whereas the present method provides for "a semi-array" containing cDNA fragments characterizing a specific "situation" for a specific cell type, according to Figure 8.

Such a surface can *i.a.* be used for screening for genes within a gene family. The "array chip" is provided and thereafter a labelled probe (which is a representative of a gene family) is allowed to hybridize to the chip under low stringency *i.e.* under conditions as described at pages 94-106 in "Nucleic acid hybridisation. A practical approach" edited by BD Hames & S J Higgins, IRL Press. A number of fragments coupled to the chip will hybridize to the probe, and these fragments can subsequently be identified, isolated and sequenced/characterized in order to determine whether they are representatives of the same gene family.

Another use of such "semi-arrays" is for determining the difference in expression pattern between a first cell or type of cells and a second cell or type of cells, the method comprising providing samples of labelled RNA or cDNA from the first and second cells or cell types and subsequently contacting



each of these samples with a chip surface as described above, and subsequently detecting the amount and distribution of bound labelled RNA or cDNA from each sample.

Under all circumstances, the chip surface with the cDNA bound thereto can e.g. be produced by the methods described in EP-0 654 061.

Yet another part of the invention pertains to a method for screening for interactions between a pre-selected protein and a polypeptide fragment, the method comprising preparing a sub-divided library of amplified cDNA fragments resulting from step e), optionally adapting the terminals of the members of the library so as to facilitate insertion in a vector, inserting the fragments into vectors, transforming a population of suitable host cells with the vectors, culturing the host cells under conditions which enable expression of correctly inserted cDNA fragments by the host cell, and subsequently assaying polypeptide fragments encoded by the inserted cDNA fragments for interaction with the pre-selected protein.

One convenient way of achieving this is by way of a two-hybrid technique, wherein the host cells are eukaryotic cells (such as fungal cells, especially yeast cells) which are mated or transfected with nucleic acid material encoding the pre-selected protein, successful mating/transfection of the cell(s) resulting in a cell or cells wherein the interaction between the pre-selected protein and a polypeptide fragment gives rise to a detectable signal.

Such methods have recently attracted a great deal of attention, i.a. as a consequence of the disclosure in Fromont-Racine et. al., Nature Genetics 16, 277-282 (1997), which is incorporated by reference herein.

One convenient system for providing the detectable signal is by use of Green Fluorescent Protein, disclosed in EP-A-0 569



170, wherein changes in fluorescent spectrum due to interactions are used as reporter.

Finally, the invention pertains to a composition for use in reverse transcription of RNA, the composition comprising

- 5 a) a first enzyme having reverse transcriptase activity at temperatures not exceeding 55°C
- b) a second enzyme having reverse transcriptase activity at elevated temperatures in the range of 45°C - 95°C (and especially the temperatures discussed above for performing reverse transcription at elevated temperatures),

10 said second enzyme having a substantially higher activity than said first enzyme in catalyzing reverse transcription at said elevated temperatures. It is preferred that the first enzyme has a substantially higher activity than said second enzyme in catalyzing reverse transcription at said temperatures not exceeding 55°C, and it is also preferred that the second enzyme has a substantially higher activity than said first enzyme in catalyzing reverse transcription at said temperatures exceeding 45°C.

## 20 DESCRIPTION OF THE PREFERRED EMBODIMENTS

First, the drawing will be briefly described.

Fig. 1

Basis of Display Of Differentially Expressed Transcripts.

Fig. 2

25 Anchor and PCR primer design.

Fig. 3

An autoradiogram of a DODET gel using the cellular set-up described in Example 1; rat pheochromocytoma PC12 cells were stimulated with the Nerve Growth Factor (NGF) and Epidermal growth factor (EGF).

30

Lanes 1 - 24, reverse transcription using the anchored poly T primer 5'-T<sub>25</sub>AA-3'

Lanes 25 - 48, reverse transcription using the anchored poly T primer 5'-T<sub>25</sub>GC-3'

- 5 Lanes 1, 5, 9, 13, 17, 21, 25, 29, 33, 37, 41, 45 represent the PC12 cells not treated.

Lanes 2, 6, 10, 14, 18, 22, 26, 30, 34, 38, 42 and 46 represent the PC12 cells treated with the NGF factor for 60 minutes.

- 10 Lanes 3, 7, 11, 15, 19, 23, 27, 31, 35, 39, 43 and 47 represent the PC12 cells treated with the NGF factor for 90 minutes

- 15 Lanes 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44 and 48 represent the PC12 cells treated with the EGF factor for 90 minutes.

Lanes 1 - 48, using the following pairs for the pre-PCR amplifications:

TaqI pre-amplification primer: 5'-CAGCATGAGTCCTGACCGA

BclI pre-amplification primer: 5'-CTCGTAGACTGCGTACCGATCA

- 20 For the second PCR amplification the following primer pairs were used:

Lanes 1-4

5'-CATGAGTCCTGACCGAA

5'-GACTGCGTACCGATCAA (5' end labelling)

- 25 Lanes 5 - 8

5'-CATGAGTCCTGACCGAA

5'-GACTGCGTACCGATCAC (5' end labelling)

Lanes 9 - 12

5'-CATGAGTCCTGACCGAA

- 30 5'-GACTGCGTACCGATCAG (5' end labelling)

Lanes 13 - 16

5' - CATGAGTCCTGACCGAA

5' - GACTGCGTACCGATCAT (5' end labelling)

Lanes 17 - 20

5 5' - CATGAGTCCTGACCGAC

5' - GACTGCGTACCGATCAA (5' end labelling)

Lanes 21 - 24

5' - CATGAGTCCTGACCGAC

5' - GACTGCGTACCGATCAC (5' end labelling)

10 Lanes 25 - 48

Repeated primer combination from lanes 1 - 24

Fig. 4a

Northern Blot of RDF01 sequence from cellular total RNA.

15 a) PC12 cells not treated, b) NGF treatment for 60 minutes,  
c) NGF treatment for 90 minutes, and d) EGF treatment for 90  
minutes.

Fig. 4b

20 Loading control, RNA extracts were electrophoresed on a 1.2%  
agarose gel containing ethidium bromide, used as a control to  
determine the relative concentration of RNA in each lane, a,  
b, c, d same as in Figure 4a

Fig. 4c

Northern Blot of RDF02 sequence from cellular total RNA.

25 a) PC12 cells not treated, b) NGF treatment for 60 minutes,  
c) NGF treatment for 90 minutes, and d) EGF treatment for 90  
minutes.

Fig. 4d

30 Loading control, RNA extracts were electrophoresed on a 1.2%  
agarose gel containing ethidium bromide, used as a control to  
determine the relative amount of RNA in each lane, a, b, c, d  
same as in Figure 4c

Fig. 5

Searching for genes modulated by a growth factor.

	Lane 1	Size marker in bp (150 bp, 200 bp, 250 bp).
	Lanes 2-6	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAA
5	Lanes 7-11	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAC
	Lanes 12-16	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAG
10	Lanes 17-21	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAT
	Lanes 22-26	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GCA

In lane 11 a downregulation is observed after 6 days treatment, whereas in lane 16 an upregulation is observed after 6 days treatment. Both modulations are due to the growth factor, since regulation is seen only when the active growth factor is present.

Fig 6

20 Searching for genes involved in bacterial resistance.

	Lane 1:	Size marker in bp (150 bp, 200 bp, 250 bp, 300 bp).
	Lanes 2-5	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAA
25	Lanes 6-9	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAC
	Lanes 10-13	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAG
	Lanes 14-17	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GAT
30	Lanes 18-21	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GCA
	Lanes 22-25	Amplification primer 5'-Com-N <sub>n1</sub> -3' where N <sub>n1</sub> is GCC

35 In lanes 8-9 a downregulation is observed, in lanes 20-21 an upregulation is observed. Both gene modulations are potential

genes involved in the resistance to the Bacteriamycin, Inosin.

Fig. 7

Principle of the technology used in Examples 4 and 5.

- 5 After ds-cDNA synthesis the DNA is digested with one 4 base pair endonuclease and anchors are ligated to the ds-cDNA ends. Using special design primers the expression profiles are obtained by amplifying the mRNAs in different expression windows (sub-fractions). The number of expression windows  
10 depends on the complexity of the sample i.e. 64 expression windows in eukaryotic.

Fig. 8

Principles of a gene discovery DNA surface (a DNA chip).

- After size separation of the DNA fragments, the DNA fragments  
15 are transferred to a nylon membrane using an electrophoretic principle. The membrane is hybridized with a complex DNA probe generated using the principle of the invention. Alternatively the membrane can be hybridized with one single gene to identify new members of a particular gene family. The  
20 membrane are in the x coordinates separated in 64 expression windows, and in the y coordinates separated in base pair size (from 50 base pair to 1200 base pair) according to principle described in figure 7.

Fig. 9

- 25 Principle of generation 64 pools of 3' END cDNAs

#### Step 1

Production of single stranded cDNA using 5'-con<sub>1</sub>-T<sub>n</sub>V oligonucleotide where con<sub>1</sub> is an oligonucleotide between 1-100 nucleotide, n is between 5-40 and V is a mixture of A, C and G.

## Step 2

Double stranded cDNA synthesis are produced using 5'Con<sub>2</sub>N<sub>x</sub> where con<sub>2</sub> is an oligonucleotide between 1-100 nucleotide, x is between 1-10, and N is a mixture of A, C, T and G. The ds-cDNA synthesis is synthesized by Klenow enzyme with the above-described oligonucleotide.

## Step 3

Pre-amplification of double stranded cDNA to amplify the double stranded cDNA, the cDNA is PCR amplified using a combination of con<sub>1</sub> and con<sub>2</sub> primers.

## Step 4

The pre-amplified cDNA is further amplified and separated in 64 pools using a combination of a labeled con<sub>1</sub> and 64 con<sub>2</sub>NNN primers in a PCR amplification procedure, where NNN are combined in 64 different ways using the nucleotides A, T, G and C.

## Step 5

Each of the 64 pools is separated using the Page electrophoresis principle.

## 20 EXAMPLES

In order to verify the functionality of the invention, examples are described below in which a developmental eukaryotic cellular system, pheochromocytoma PC12, was employed.

Nerve Growth Factor (NGF) induces growth arrest and neurone outgrowth in the *in vitro* PC12 cell system. Other growth factors, such as epidermal growth factor (EGF), support survival and stimulate growth. NGF-induced genes, include the immediate early genes, which encode transcription factors, such as c-fos and c-myc. The products of the immediate early genes are thought to be involved in regulating the expression of genes, associated with the neuronal phenotype for example neurofilaments, peripherin, GAP43 and transin.

In order to identify new early genes involved in neuronal differentiation and proliferation, the following DODET method is used for identify such genes.

5 In the following examples, it is demonstrated how efficiently the method of the invention can be applied to such cellular systems.

#### EXAMPLE 1

10 The rat pheochromocytoma PC12 cells were grown (*in vitro*) in the presence and absence of Nerve Growth Factor (NGF) and epidermal growth factor (EGF) under growth conditions described elsewhere (Saltiel et al. 1996).

The total RNA was isolated using the standard single-step method by Chomczynski and Sacchi according to Sambrook et al 1989.

15 Total RNA concentration was determined spectrophotometrically and then adjusted to 0.2  $\mu\text{g}/\mu\text{l}$ . This RNA was used directly in the Northern analysis.

20 For DODET 4 x 0.5  $\mu\text{g}$  total RNA was reverse transcribed in separated pools using the primer 5'-T<sub>25</sub>AA-3'. The same procedure was performed using the 5'-T<sub>25</sub>GC-3' poly-dT anchored primers, giving a total of 2 x 4 x 0.5  $\mu\text{g}$  of RNA.

#### First strand synthesis

20.0 $\mu\text{l}$	total RNA amount between 0.3 to 1.0 $\mu\text{g}$ RNA
3.0 $\mu\text{l}$	5'-T <sub>25</sub> AA-3' Conc. 100 ng/ $\mu\text{l}$ or 5'-T <sub>25</sub> GC-3' Conc.
25	100 ng/ $\mu\text{l}$
5.0 $\mu\text{l}$	10 x cDNA buffer (buffer B from Epicentre Technologies # R19250)
2.0 $\mu\text{l}$	dNTPs (25 mM from Pharmacia Biotech)
1.0 $\mu\text{l}$	SuperScript II RT (200 U/ $\mu\text{l}$ ) (Gibco BRL # 18064-
30	014)



5.0  $\mu$ l Retrotherm RT (1 U/ $\mu$ l) (Epicentre Technologies  
#R19250)  
14.0  $\mu$ l H<sub>2</sub>O

To obtain high specificity, the cDNA reaction was incubated  
5 at 50°C for 30 minutes followed by 1 hour incubation at 70°C.

#### Second strand synthesis:

To the first strand reaction, add the following components

15.0  $\mu$ l 10 x cDNA buffer  
3.0  $\mu$ l Hybridase Thermostable RNase (1 U/ $\mu$ l) (Epicentre  
10 Technologies #H39050)  
1.0  $\mu$ l rBst thermostable DNA polymerase (1 U/ $\mu$ l) (Epi-  
centre Technologies #BH1100)  
81.0  $\mu$ l H<sub>2</sub>O

Incubate at 65°C for 1 hour.

15 The resulting double stranded cDNA was phenol extracted and  
precipitated and resuspended in 20  $\mu$ l of H<sub>2</sub>O. Half of this  
volume was checked on gel; if a smear between 100 bp and 3000  
bp was observed, the rest of the cDNA was used for DODET  
template production. The resulting cDNAs were digested with  
20 10 U of each of the thermostable restriction enzymes *Taq*I and  
*Bcl*II at 50°C for 2 hours. To this mixture, DODET adapters  
were added and ligated to the ends of the restriction frag-  
ments with T<sub>4</sub> DNA ligase (1U) resulting in the primary tem-  
plate. 8-15 cycles of non-radioactive pre-amplification,  
25 using primers complementary to the DODET adapters, were  
performed on a small aliquot (1/10<sup>th</sup> volume) of the primary  
template (94°C denaturation; 30 s, 56°C annealing; 30 s, 72°C  
polymerisation; 1 min). The products of the amplification  
(termed secondary template) were also checked on a 1.5%  
30 agarose gel. As expected, fragment sizes were predominantly  
between 100 bp and 1000 bp. All amplification reactions were  
carried out on a PE-9600 thermocycler using *Taq* DNA-polyme-

rase, both from Perkin Elmer Corp. (Norwalk, CT, USA). The final template was then diluted 10 fold with H<sub>2</sub>O.

The adapters ligated to the restriction fragments, the pre-amplification and active PCR are given below:

- 5    *TaqI* adapter:            5'-CAGCATGAGTCCTGAC  
                                 TACTCAGGACTGGC-5'

*TaqI* pre-amplification primer: 5'-CAGCATGAGTCCTGACCGA

*TaqI* amplification primer:        5'-CATGAGTCCTGACCGAN  
(N = A or C or G or T)

- 10   *BclI* adapter:        5'-CTCGTAGACTGCGTACC  
                                 CTGACGCATGGCTAG-5'

*BclI* pre-amplification primer: 5'-CTCGTAGACTGCGTACCGATCA

*BclI* amplification primer: 5'-GACTGCGTACCGATCAN  
(N = A or C or G or T)

- 15   For PCR all the different combinations of one extension  
      (denoted as N above) were available, giving a total of 4<sup>2</sup>  
      primer combinations. All oligonucleotides were obtained from  
      DNA Technology (Aarhus, Denmark).

- 20   Radioactive labelling of the *BclI* primer was performed using  
      1U of T<sub>4</sub> polynucleotide kinase. Thermocycling was carried out  
      essentially as described above but with 35 cycles and includ-  
      ing an 11 cycle touch-down (the annealing temperature was  
      reduced from 65°C to 56°C in 0.7°C steps for 11 cycles and  
      subsequently maintained at 56°C for 23 cycles). Samples were  
25   then boiled after the addition of dye and 50% formamide and  
      separated on a 5% polyacrylamide sequencing type gel (GIBCO  
      BRL Life Technologies Inc., Gaithersburg, MD, USA). All gels  
      were run at standard conditions, such that the 70 bp marker  
      was 3 cm from the bottom of the gel, giving good resolution

between 70-800 bp. Gels were then dried directly onto Whatman 3M paper on a slab gel dryer. Labelled DNA fragments were visualised by autoradiography. Gels and films were positionally marked prior to development. The 1 base selective extensions were chosen empirically to yield approximately 50 radioactively labelled fragments per lane.

Bands, identified on the autoradiogram as interesting, were lined up with markings on the film and the dehydrated gel and were excised. Excised fragments were monitored for activity. The gel fragments were isolated using GENE CLEAN (BIO101, California USA). DNA was then recovered according to the manufacturer's recommendations. DNA fragments could then be reamplified using the same PCR conditions and primers as used in the initial PCR; however, 15 cycles generally yielded sufficient product for cloning. Cloning was achieved using unpurified PCR product and the vector display-p123T (Display Systems Biotech, USA). Conditions were used as recommended by the manufacturer.

#### EXAMPLE 2

Figure 3 shows a typical DODET gel produced by amplification of template derived from treatment of PC12 cells with NGF or EGF.

Total RNA was reverse transcribed with the 5'-T<sub>25</sub>AA-3' and T<sub>25</sub>GC-3' poly-dT anchored primers, and after anchor ligation pre-amplification with *Bcl*I and *Taq*I pre-amplification primer pairs was performed.

6 out of 16 possible primer combinations are shown, using 1 selective base, at each restriction enzyme site (Figure 3). The largest visible products (Figure 3) are approximately 1000 bp in size and the lower end of the gel corresponds to approximate 100 bp. In this size window an average of 50 bands can be scored for each primer combination. In Figure 3, various expression patterns can be detected.

Due primarily to the stringent conditions possible in DODET, resolution of the banding pattern is high while the level of background remains at acceptable levels (Figure 3). Furthermore, quite radical changes in the intensity of individual  
5 bands over the treatment period do not seem to affect the patterns of other bands in the same lane.

It is, therefore, possible to conclude that the PCR remains proportionally independent on the concentration of individual substrates in the reaction.

10 The use of an optimised combination of standard protocols described above for isolating, re-amplifying and cloning individual RDFs, has allowed the identification of a number of transcripts associated with differentiation and proliferation events.

15 Four RDFs were isolated for further analysis, Figure 3, bands a, b, c and d.

Sequence analysis revealed that RDF a = RDF b and RDF c = RDF d, as illustrated in Figure 3.

In all cases appropriate terminal sequences with the correct  
20 1 selective base extensions used in the PCR could be retrieved, demonstrating the stringency and fidelity of the system (data not shown).

### EXAMPLE 3

During scanning of the PC12 cellular systems treated with NGF  
25 or EGF with different primer combinations, two RDFs (designated RDF01 and RDF02) exhibiting a differential expression during the NGF treatment were isolated (RDF01 = RDF a = RDF b and RDF02 = RDFc = RDF d. in Figure 3).

After re-amplification, sub-cloning and DNA sequencing,  
30 further DNA analysis revealed two unknown RDFs upregulated

after 60 minutes NGF treatment or 90 minutes EGF treatment in the PC12 cellular system. The nucleotide sequences of both RDF01 and RDF02 show less than 10% homology to any existing gene in the GeneBank or EMBL databases.

- 5 The expression of RDF01 and RDF02 was further analyzed using Northern blot (Figure 4a - 4d). Here, transcripts could clearly be detected at 60 minutes NGF treatment or 90 minutes EGF treatment of the PC12 cells, confirming the results obtained using the DODET method, as illustrated in Figure 3.
- 10 Experiments to clone the full length of RDF01 and RDF02, and biological characterisation of their involvement in the differentiation and proliferation, of the PC12 cellular system are currently under investigation.

#### EXAMPLE 4

- 15 *Searching for genes modulated by a growth factor.*

A human cell line was treated with a growth factor and RNA was isolated at various time points as indicated below.

- |    |   |
|----|---|
| 1  | Cell without any treatment (lanes 2, 7, 12, 17, 22)     |
| 2  | Cell treated with helper agent, 1 day (lanes 3, 8, 13,  |
| 20 | 18, 23)   |
| 3  | Cell treated with helper agent and growth factor, 1 day |
|    | (lanes 4, 9, 14, 19, 24)                                |
| 4  | Cell treated with helper agent, 6 days (lanes 5, 10,    |
|    | 15, 20, 25)   |
| 25 | 5 Cell treated with helper agent and growth factor, 6   |
|    | days (lanes 6, 11, 16, 21, 26)                          |

Human RNA was isolated and the gene discovery analysis was performed essentially as described in the legend to Fig. 3. 5 out of 64 amplification primers are shown in Fig. 5, each covering a certain portion of the mRNA pool in the human cell

30 line. The expression analysis was performed on an ALFexpress,

an automated fragment analyzer from Pharmacia Biotech, using a Cy5 label.

In lane 11 a downregulation is observed after 6 days of treatment. In lane 16 an upregulation is observed after 6 days of treatment. Both modulations are due to the growth factor, since regulation is only observed with the active growth factor present.

#### EXAMPLE 5

*Searching for genes involved in bacterial resistance to antibiotics.*

A *Listeria monocytogenes* strain was treated with the Bacteriamycin, Inosin. RNA from a strain resistant to Inosin was further investigated.

1. Bacterial clone 1 without any treatment (lanes 2, 6, 10, 14, 18)
2. Bacterial clone 2 without any treatment (lanes 3, 7, 11, 15, 19)
3. Bacterial clone 3 resistant to Inosin (lanes 4, 8, 12, 16, 20)
4. Bacterial clone 4 resistant to Inosin (lanes 5, 9, 13, 17, 21)

Bacterial RNA was isolated by standard techniques and the gene discovery analysis was performed according to Example 4 and Fig. 5, with the exception that a 5'-NNNNNNYYA primer was used for first strand synthesis.

6 of 64 amplification primers are shown in Figure 6, each covering a certain portion of the mRNA pool in the prokaryotic cell system. The expression analysis was performed on an ALFexpress, an automated fragment analyzer from Pharmacia Biotech, using a Cy5 label.




In lanes 8 and 9 a downregulation is observed and in lanes 20 and 21 an upregulation is observed. Both gene modulations are potential genes involved in the resistance to Inosin.

## CLAIMS


1. A method for preparing a normalized sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample, the method comprising the steps of

- 5 a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer having the general formula



10 wherein  $\text{Con}_1$  is any sequence between 1-100 nucleotides, dT is deoxythymidynyl, V is A, G or C, N is A, G, C or T,  $n2$  is an integer  $\geq 1$ ,  $n3$  is 0 or 1, if  $n3$  is 0 then  $n4$  is 0, and if  $n3$  is 1  $n4$  is an integer  $\geq 0$ , thereby obtaining first strand cDNA fragments,

- 15 b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, and a second cDNA primer with the general formula



20 wherein  $\text{Con}_2$  is any sequence between 1-100 nucleotides and can be different or identical to  $\text{con}_1$ ,  $\text{N}_x$  is A, G, T or C, and  $x$  is an integer  $\geq 0$ , in a appropriate enzyme/-buffer solution which comprises the DNA pol I enzyme or the Klenow fragment of the DNA pol I enzyme, all four deoxyribonucleoside triphosphates and standard buffer and  
25 temperature conditions, thereby obtaining double stranded cDNA fragments, and

- 30 c) subjecting the cDNA fragments obtained in step b) to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used a set of amplification primers having the general formula

$$5' - \text{Con}_3 - \text{N}_{n1} - 3'$$

I

5 wherein  $\text{Con}_3$  is a sequence identical to either  $\text{Con}_1$  or  $\text{Con}_2$  or both, N is A, G, T or C, and  $n1$  is an integer  $\geq 0$ , wherein at least one set of primers has the general formula I where  $n > 0$ , said at least one set being capable of priming amplification of any nucleotide sequence complementary in its 5'-end to  $\text{Con}_1$  or  $\text{Con}_2$ .

10 2. A method for preparing a normalized sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample, the method comprising the steps of

- a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer, thereby obtaining first strand cDNA fragments,
- 15 b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, thereby obtaining double stranded cDNA fragments,
- c) digesting the double stranded cDNA fragments with at least one restriction endonuclease, thereby obtaining  
20 cleaved cDNA fragments,
- d) ligating at least two adapter fragments to the cleaved cDNA fragments obtained in step c), so as to obtain ligated cDNA fragments, and
- 25 e) subjecting the ligated cDNA fragments obtained in step d) to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used, for an adapter fragment used in step d), a set of amplification primers having the general formula

$$5' - \text{Com} - \text{N}_{n1} - 3'$$

II

wherein Com is a sequence complementary to at least the 5'-end of an adapter fragment which is ligated to the 3'-end of a cleaved cDNA fragment, N is A, G, T, or C, and n1 is an integer  $\geq 0$ , and wherein at least one set of  
5 primers has the general formula I where  $n1 > 0$ , said at least one set being capable of priming amplification of any nucleotide sequence ligated in its 3'-end to the adapter fragment complementary in its 5'-end to Com.

3. A method according to claim 1 or 2, wherein the mRNA is of  
10 eukaryotic, Archae or prokaryotic origin.

4. A method according to any of claims 1-3, wherein the reverse transcription is performed under high stringency conditions.

5. A method according to any of the preceding claims, wherein  
15 the reverse transcription is carried out at a temperature in the range from about 45°C to about 95°C by use of an enzyme having reverse transcriptase activity at said temperature.

6. A method according to claim 5, wherein the enzyme is thermostable, such as an enzyme selected from the group  
20 consisting of a DNA polymerase with reverse transcriptase activity derived from thermophilic eubacteria, such as Taq (Thermus aquaticus), Stoffel (Thermus aquaticus), Tht (Thermus thermophilus), Tfl/Tub (Thermus flavus), Tru (Thermus Ruber), Tca (Thermus caldophilus), Tfil (Thermus filiformis),  
25 Tbr (Thermus Brockianus), Bst (B. Stearothermophilus), Bca (B. Caldotenax YT-G), Bcav (B. Caldovelox YT-F), FjSS3-B.1 (Thermotoga FjSS3-B.1), Tma (Thermus Maritima), UITma (T. Maritima), Tli (T. Litoralis), Tli exo- (T. Litoralis), 9°N-7 (Thermococcus sp.), BG-D (Pyrococcus sp.), Pfu (P. furiosus),  
30 Pwo (P. woesei), Sac (S. Acidocaldarius), SsoI (S. Solfataricus), Tac (T. Acidophilum), and Mth (Methananococcus Voltae).

7. A method according to any of claims 1-4, wherein the reverse transcription is carried out at a temperature in the

range from about 25°C to about 55°C by use of an enzyme having reverse transcriptase activity at said temperature.

8. A method according to claim 7, wherein the enzyme is a reverse transcriptase, such as a reverse transcriptase selected from reverse transcriptase from AMV (Avian Myeloblastosis Virus), M-MuLV (murine M-MuLV pol gene), or HIV-1 (HIV virus).

9. A method according to any of claims 1-4, wherein the reverse transcription is carried out in two subsequent steps, the first step comprising carrying out reverse transcription as defined in claim 7 or 8, and the second step comprising carrying out reverse transcription as defined in claim 5 or 6.

10. A method according to claim 9, wherein reverse transcription in the two steps is effected by non-identical enzymes having reverse transcriptase activity.

11. A method according to claim 10, wherein the non-identical enzymes are added separately in each step or are present in both steps.

12. A method according to claim 11, wherein the activity of the enzyme which is active in the first step is substantially abolished in the second step.

13. A method according to any of claims 9-12, wherein the enzyme effecting reverse transcription in the first step is reverse transcriptase from MMuLV, AMV or HIV-1 and/or the enzyme effecting reverse transcription in the second step is Tth or Taq.

14. A method according to any of the preceding claims, wherein the at least one cDNA primer includes an oligo or poly dT tail in the 3' end.

15. A method according to claim 14, wherein the at least one cDNA primer has the general formula  $5'-dT_{n2}-V_{n3}-N_{n4}-3'$ , wherein dT is deoxythymidiny1, V is A, G, or C, N is A, G, C or T,  $n2$  is an integer  $\geq 1$ ,  $n3$  is 0 or 1, if  $n3$  is 0 then  $n4$  is 0, and if  $n3$  is 1 then  $n4$  is an integer  $\geq 0$ .
16. A method according to claim 15, wherein, when  $n3$  is 1,  $3 \times 4^{n4}$  groups of cDNA primers are used, each group being distinct from any one of the other groups with respect to the structure  $-V_{n3}-N_{n4}-$ .
17. A method according to claim 16, wherein the pool of mRNA is subdivided into  $3 \times 4^{n4}$  aliquots which are each subjected separately to step a) utilising one of the  $3 \times 4^{n4}$  groups of cDNA primers, thereby obtaining a subdivision of the first strand cDNA into  $3 \times 4^{n4}$  separate pools.
18. A method according to claim 16 or 17, wherein  $n4$  is 0 or 1.
19. A method according to any of claims 1-13, wherein the at least one cDNA primer does not include a poly or oligo dT tail in the 5'-end, or wherein at least two cDNA primers are used of which at least one includes a poly or oligo dT tail in the 5'-end and of which at least one second does not include a poly or oligo dT tail in the 5'-end.
20. A method according to claim 19, wherein the cDNA primer which does not include a poly or oligo dT tail in the 5'-end has the following structure
- $$5'-N_xTTA-3' \text{ or } 5'-N_xCTA-3' \text{ or } 5'-N_xTCA-3',$$
- wherein N is A, G, T, or C, and x is an integer  $1 \leq x \leq 20$ .
21. A method according to any of the preceding claims, wherein step b) is carried out under conditions which minimize the



formation of mismatches between nucleotides in the first and second cDNA strands.

22. A method according to any of the preceding claims, wherein the at least one restriction enzyme is chosen so as to ensure that at least 60% of cDNA's are cleaved.

23. A method according to any of the preceding claims comprising the use of at least one restriction endonuclease which upon cleavage of cDNA results in cleaved cDNA fragments having sticky ends.

24. A method according to any of the preceding claims, wherein the at least one restriction enzyme is chosen so as to cleave each complete cDNA into an average of about 3 fragments.

25. A method according to any of the preceding claims, comprising the use of a rare 4 base cutter as at least one restriction endonuclease, such as the 4 base cutter *AccI*, *AluI*, *BfaI*, *BstUI*, *Csp6I*, *DpnI*, *DpnII*, *HaeIII*, *HhaI*, *HinPII*, *HpaII*, *MboI*, *MnlI*, *MseI*, *MspI*, *NlaIII*, *RsaI*, *Sau3AI*, *TaiI*, *TaqI*, and *Tsp509I*.

26. A method according to any of the preceding claims, wherein one restriction enzyme is used.

27. A method according to any of claims 1-21, which comprises the use of a first restriction enzyme which statistically cleaves at least 20% of complete cDNA derived from the mRNA sample into two subfragments, and of a second restriction enzyme which statistically cleaves at least 50% of said subfragments into 3 further subfragments.

28. A method according to any of the preceding claims, wherein, in step d), at least one termination fragment is also ligated to the 3'-end of single strands of cleaved cDNA fragments, said at least one termination fragment introducing a

block against DNA polymerization in the 5'→3' direction  
setting out from the at least one termination fragment and  
said at least one termination fragment being unable to anneal  
to any primer of the at least two primer sets in step e)  
5 during the molecular amplification procedure.

29. A method according to claim 28, wherein the at least one  
termination fragment comprises or is a chemically modified  
nucleotide sequence.

30. A method according to claim 29, wherein the chemically  
10 modified nucleotide sequence comprises a dideoxynucleotide in  
the 3'-end.

31. A method according to claim 30, wherein the dideoxy-  
nucleotide is covalently attached to the nucleotide strand.

32. A method according to any of the preceding claims, where-  
15 in the ligation of adapter and/or termination fragments to  
the cleaved cDNA fragments in step d) is achieved by anneal-  
ing the adapter fragments to sticky ends of the cDNA result-  
ing from the cleavage in step c) and subjecting the product  
to the action of an enzyme having DNA ligase activity.

20 33. A method according to any of the preceding claims, where-  
in the at least one set of amplification primers of formula I  
or II wherein  $n_1$  is  $\geq 0$  has  $n_1=1$ ,  $n_1=2$ ,  $n_1=3$ , or  $n_1=4$ .

34. A method according to any of the preceding claims, where-  
in  $n_1=0$  in the at least one set of amplification primers  
25 having formula I or II.

35. A method according to claim 34, wherein the set of ampli-  
fication primers having  $n_1=0$  in formula I or II is labelled.

36. A method according to any of the preceding claims, where-  
in the set of amplification primers having formula I or II

wherein  $n_1 > 0$  comprises all possible combinations and permutations of A, G, T and C in the group  $N_{n_1}$ .

37. A method according to any of the preceding claims, wherein the ligated cDNA fragments are sub-divided into a number  
5 of pools prior to the molecular amplification in step e), each pool being subjected to the amplification using a subset of the set of amplification primers.

38. A method according to claim 36, wherein the subset of  
10 amplification primers used for each pool comprises a primer as defined in claim 35.

39. A method according to any of the preceding claims, which comprises the use of one amplification primer as defined in claim 35, and of one set of primers as defined in claim 36.

40. A method according to claim 39, wherein the ligated cDNA  
15 fragments of step d) are subdivided into  $4^{n_1}$  pools which are each subjected separately to step e) wherein is used one amplification primer as defined in claim 35 and one primer from the set of amplification primers as defined in claim 36, said one primer being distinct from any one of the primers  
20 used for amplifying any of the other pools.

41. A method according to any of the preceding claims, which comprises the further step of separating amplified fragments obtained from the molecular amplification procedure.

42. A method according to claim 41, wherein the separation is  
25 performed by gel electrophoresis or chromatography.

43. A method according to claim 41 or 42, which further comprises the step of identifying separated amplified fragments.

44. A method according to claim 43, wherein the identifica-  
30 tion is performed by visualization.

45. A method according to claim 44, wherein labelled nucleotides are visualized, the labelled nucleotides being part of a probe or of the amplified fragments.

46. A method according to claim 45, wherein the labelled  
5 nucleotides are the labelled nucleotides being part of the labelled primers as defined in claim 35.

47. A method according to claim 46, wherein the visualization is performed by incorporating radioactive or fluorescent  
10 alpha dNTP into the cDNA fragment during PCR, where N = A, C, T, U or G.

48. A method for determining the presence of an expression product in a cell or group of cells, the method comprising providing an RNA-containing sample from the cell or group of cells and subjecting the sample to the method according to  
15 any of claims 1-47, and thereafter performing a comparison of the thus identified amplified cDNA fragments with a database output, said database output comprising a computer-generated list of molecular weights of restriction DNA fragments of known sequences, said list being prepared by

- 20 - inputting and storing DNA sequence data in a database as virtual DNA sequences,
- subsequently simulating cleavage of the virtual DNA sequences with the at least one restriction nuclease and storing the resulting simulated cleavage products as  
25 virtually cleaved DNA fragments,
- simulating ligation to the virtually cleaved fragments of the at least two adapter fragments and storing the results as virtually ligated DNA fragments,
- for each individual combination of primers used in step  
30 e), grouping the virtually ligated DNA fragments susceptible to amplification by said combination of primers in the same group,

- determining, in each group, the absolute and/or relative molecular weight of each virtually ligated DNA fragment, and
- outputting the content of each group in the form of a list comprising the absolute and/or relative molecular weights of the virtually ligated fragments in the group.

49. A method according to claim 48, wherein the input DNA sequence data are linked to data relating to the genetic origin of the DNA sequence data and optionally to data relating to functional features relating to the genetic origin.

50. A method according to claim 48, wherein the output indication further comprises information about the genetic origin of the virtually ligated DNA fragment and optionally information about functional features associated with the genetic origin.

51. A method according to any of claims 38-50, wherein the comparison is performed by inputting the identified amplified cDNA fragments in a format which allows automated comparison with the database output, or, alternatively, by outputting the database output in a format which allows for direct comparison between the separated amplified cDNA fragments and the database output.

52. A method for determining change in expression, compared to the expression in a reference cell or reference group of cells, of an expression product in a cell or group of cells which has been subjected to a first set of conditions influencing the expression pattern of said cell or group of cells, said reference cell or group of cells being subjected to a second set of conditions, the method comprising providing an RNA-containing sample from the cell or group of cells and subjecting the sample to the method according to any of claims 43-47 thereby obtaining data describing the amplified cDNA fragments derived from the sample, providing reference data describing amplified cDNA fragments derived

from an RNA-containing reference sample from the reference cell or reference group of cells, the reference data being obtained by having previously subjected the reference sample to the method according to any of claims 43-47,  
5 subsequently performing a comparison of the data and the reference data to identify those cDNA fragments which are expressed at different levels in the two data sets, and thereafter using the differentially expressed cDNA fragments to determine which expression products are subject to a  
10 change in expression level.

53. A method according to claim 52, wherein the data and reference data are selected from the group consisting of the apparent molecular weights of the amplified DNA fragments, the  $M_r$  of the amplified DNA fragments, the absolute amount of  
15 the amplified DNA fragments, and the relative amounts of the amplified DNA fragments.

54. A method according to claim 52, wherein the reference data are extracted from a database containing the reference data defined in claim 53 and optionally further information  
20 relating to the genetic origin of each amplified cDNA fragment from the reference.

55. A method for diagnosing a disease in a subject, said disease being characterized by a deviating expression level of at least one expression product in at least one cell type,  
25 the method comprising providing an RNA-containing sample derived from the at least one cell type, subjecting the sample to the method according to any of claims 43-47 thereby obtaining data describing the amplified cDNA fragments derived from the sample, providing reference data describing  
30 amplified cDNA fragments derived from a RNA-containing reference sample derived from the same type of cell from a subject not suffering from the disease, the reference data being obtained by having previously subjected the reference sample to the method according to any of claims 43-47, and



subsequently performing a comparison of the data and the reference data with respect to those cDNA fragments which are known to be related to the disease, and assessing whether a significant difference in the data and reference data exists  
5 so as to establish whether the expression level of the expression product deviates or not.

56. A method according to claim 55, wherein the data and reference data are selected from the group consisting of the apparent molecular weights of the amplified DNA fragments,  
10 the  $M_r$  of the amplified DNA fragments, the absolute amount of the amplified DNA fragments, and the relative amounts of the amplified DNA fragments.

57. A method according to claim 55, wherein the reference data are extracted from a database containing the reference  
15 data defined in claim 56 and optionally further information relating to the genetic origin of each amplified cDNA fragment from the reference.

58. A method of synthesizing first strand cDNA, the method comprising subjecting a sample comprising mRNA to reverse  
20 transcription wherein, in a first step performed at a temperature not exceeding 55°C, a first enzyme is used having a substantial reverse transcriptase activity at said temperature not exceeding 55°C, and, in a subsequent second step performed at an elevated temperature in the range of 45°C -  
25 95°C, a second enzyme is used having a substantial reverse transcriptase activity at said elevated temperature, said first enzyme being substantially inactive.

59. A method according to claim 58, wherein both enzymes are present in both steps.

30 60. A method according to claim 58, wherein said first enzyme has a substantially higher activity than said second enzyme in catalyzing reverse transcription at said temperature not exceeding 55°C.

61. A method according to claim 59 or 60, wherein said second enzyme has a substantially higher activity than said first enzyme in catalyzing reverse transcription at said elevated temperature.
- 5 62. A method according to any of claims 58-61, wherein said first enzyme is selected from the group consisting of non-thermostable reverse transcriptases and wherein said second enzyme is selected from the group consisting of thermostable DNA polymerases with reverse transcriptase activities.
- 10 63. A composition for use in reverse transcription of RNA, the composition comprising
- a) a first enzyme having reverse transcriptase activity at temperatures not exceeding 55°C
- b) a second enzyme having reverse transcriptase activity
- 15 at elevated temperatures in the range of 45°C - 95°C,
- said second enzyme having a substantially higher activity than said first enzyme in catalyzing reverse transcription at said elevated temperatures.
- 20 64. A composition according to claim 63, wherein said first enzyme has a substantially higher activity than said second enzyme in catalyzing reverse transcription at said temperatures not exceeding 55°C.
- 25 65. Use of a thermostable enzyme having reverse transcriptase activity in the preparation of a composition for use in reverse transcription of RNA which has previously been *in vitro* reverse transcribed by another enzyme having reverse transcriptase activity.
66. A method of preparing a surface (chip) coated with cDNA fragments, the method comprising

- subjecting an RNA-containing sample to the method of any of claims 41-47, and
- transferring the separated amplified cDNA fragments to a chip surface adapted to stably bind the separated amplified cDNA fragments while maintaining the spatial relative distribution pattern thereof.

67. A method of preparing a surface (chip) coated with cDNA fragments, the method comprising

- subjecting an RNA-containing sample to the method of any of claims 1-40,
- separating, by electrophoresis, the thus obtained amplified cDNA fragments on a particular surface adapted to stably bind the separated amplified cDNA fragments while maintaining the relative distribution pattern after electrophoresis.

68. A method according to claim 67, wherein the electrophoresis is in the form of microelectrophoresis.

69. A method according to any of claims 66-68, wherein the transfer is accomplished by a electrophoretic blotting technique.

70. A method according to any of claims 66-68, wherein the transfer is accomplished by photo-activated organic or inorganic chemistry techniques.

71. A surface having cDNA stably bound thereto, said surface being obtainable by the method according to any of claims 66-70.

72. A method for the screening for genes within a family of genes, the method comprising providing a surface according to claim 71, wherein cDNA stably bound to the surface is hybridized under low stringency conditions to a detectably label-

led nucleic acid which is a representative of a gene family, and subsequently analyzing fragments of the chip to which hybridization has occurred so as to determine whether such fragments are related to the same gene family.


- 5 73. A method for determining the difference in expression pattern between a first cell or type of cells and a second cell or type of cells, the method comprising providing sam-  
ples of labelled RNA or cDNA from the first and second cells  
or cell types and subsequently contacting each of these  
10 samples with a surface according to claim 71, and subsequent-  
ly detecting the amount and distribution of bound labelled  
RNA or cDNA from each sample.

74. A method for screening for interactions between a pre-  
selected protein and a polypeptide fragment, the method  
15 comprising preparing a sub-divided library of amplified cDNA  
fragments according to the method of any of claims 1-47,  
optionally adapting the terminals of the members of the  
library so as to facilitate insertion into a vector, insert-  
ing the fragments into vectors, transforming a population of  
20 suitable host cells with the vectors, culturing the host  
cells under conditions which enable expression of correctly  
inserted cDNA fragments by the host cell, and subsequently  
assaying polypeptide fragments encoded by the inserted cDNA  
fragments for interaction with the pre-selected protein.

- 25 75. A method according to claim 74, wherein assaying of the  
polypeptide fragments is performed by a two-hybrid technique,  
wherein the host cells are eukaryotic cells which are mated  
or transfected with nucleic acid material encoding the pre-  
selected protein, successful mating/transfection of the  
30 cell(s) resulting in a cell or cells wherein the interaction  
between the pre-selected protein and a polypeptide fragment  
gives rise to a detectable signal.


76. A method according to claim 75, wherein a fungus, such as  
a yeast cell, is used in the mating/transfection.

77. A method according to claim 75 or 76, wherein the detectable signal is provided by Green Fluorescent Protein.

**Fig. 1 Basis of DODET**




# ● Fig. 1 Basis of DODET (continue)



3/12


● **Fig. 2 Anchor and PCR primer design**



**Continue next page**


4/12

• Fig. 2 Anchor and PCR primer design (continue)



5/12


Fig. 3



SUBSTITUTE SHEET (RULE 26)


6/12

Fig. 4a



a b c d

Fig. 4b




a b c d

SUBSTITUTE SHEET (RULE 26)


7/12

Fig. 4c



a b c d

Fig. 4d




a b c d

SUBSTITUTE SHEET (RULE 26)



8/12

Fig. 5




SUBSTITUTE SHEET (RULE 26)




9/12

Fig. 6



10/12


Fig. 7



11/12


Fig. 8

displayARRAY principle



12/12

Fig. 9



## INTERNATIONAL SEARCH REPORT

Int l Application No

PCT/DK 98/00186

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/10 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 18176 A (DANA FARBER CANCER INST INC) 16 September 1993	1, 3, 7
Y	see claims 1-62; figures 1-5	2, 48, 52, 55
Y	--- EP 0 743 367 A (MAX PLANCK GESELLSCHAFT) 20 November 1996 see page 3, line 6 - page 6, line 2 see page 6, line 19 - line 27; claims 1-26; figures 1, 3	2, 48, 52, 55
A	--- WO 97 05286 A (UNIV YALE ; WEISSMAN SHERMAN (US); PRASHAR YATINDRA (US)) 13 February 1997 see the whole document --- -/-	1-77

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 November 1998

Date of mailing of the international search report

17. 11. 1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

# INTERNATIONAL SEARCH REPORT

Int. lional Application No  
PCT/DK 98/00186

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 13369 A (SCRIPPS RESEARCH INST) 18 May 1995 see the whole document ---	1-77
A	BAUER D ET AL: "DIFFERENTIAL DISPLAY RESERSE TRANSCRIPTION PCR" BIOTEC, no. 2, April 1994, page 32, 34/35 XP000198999 see the whole document ---	1-77
A	WO 95 08647 A (UNIV COLUMBIA ;SOARES MARCELO B (US); EFSTRATIADIS ARGIRIS (US)) 30 March 1995 see the whole document ---	1-77
A	R. WILLIAMSON: "The preparation and screening of a cDNA clone bank" GENETIC ENGINEERING 1 , ED. R. WILLIAMSON, ACADEMIC PRESS, INC. ,1981, pages 1-59, XP002076424 New York, US see page 12, paragraph 2 - page 19, paragraph 1 ---	1-77
A	"Construction and analysis of cDNA libraries" MOLECULAR CLONING, SECOND EDITION, J. SAMBROOK, E.F. FRITSCH AND T. MANIATIS, , 1989, pages 8.11-8.17, XP002076425 CSH, US see the whole document ---	1-77
A	PATANJALI S R ET AL: "CONSTRUCTION OF A UNIFORM-ABUNDANCE (NORMALIZED) CDNA LIBRARY" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 88, no. 5, 1 March 1991, pages 1943-1947, XP000368687 ---	1-77
P, X	WO 97 29211 A (US HEALTH ;WEINSTEIN JOHN N (US); BOULAMWINI JOHN (US)) 14 August 1997  see page 6, line 18 - page 11, line 27; claim 1; figures 1,2 see page 19, line 33 - page 20, line 4 -----	1-3,7,8, 14-17, 19, 21-26, 32-46, 48,52

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 98/00186

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.



## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

## 1. Claims: 1-57,66-77

A method for preparing a normalized sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample, the method comprising the steps of a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer, b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, and a second primer and c) subjecting the cDNA fragments obtained in step b) to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used a set of amplification primers; a method for preparing a normalized sub-divided library of amplified cDNA fragments from the coding region of mRNA contained in a sample, the method comprising the steps of a) subjecting the mRNA derived from the sample to reverse transcription using at least one cDNA primer, b) synthesizing second strand cDNA complementary to the first strand cDNA fragments by use of the first strand DNA fragments as templates, thereby obtaining double stranded cDNA fragments, c) digesting the double stranded cDNA fragments with at least one restriction endonuclease, thereby obtaining cleaved cDNA fragments, d) ligating at least two adapter fragments to the cleaved cDNA fragments obtained in step c), so as to obtain ligated cDNA fragments, and e) subjecting the ligated cDNA fragments in step d) to a molecular amplification procedure so as to obtain amplified cDNA fragments, wherein is used, for an adapter fragment used in step d), a set of amplification primers; a method for determining the presence of an expression product in a cell or a group of cells using said method; a method for determining change in expression, compared to the expression in a reference cell or reference group of cells; a method of diagnosing a disease in a subject; a method of preparing a surface (chip) coated with cDNA fragments; a method for screening for genes within a family of genes; a method for screening for interactions between a pre-selected protein and a polypeptide fragment; said method wherein assaying of the polypeptide fragments is performed by a two-hybrid technique;

## 2. Claims: 58-65

A method of synthesizing first strand cDNA, the method comprising subjecting a sample comprising mRNA to reverse transcription wherein, in a first step performed at a temperature not exceeding 55°C, a first enzyme is used having a substantial reverse transcriptase activity at a temperature not exceeding 55°C, and, in a subsequent second step performed at an elevated temperature in the range of

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

45°C-95°C, a second enzyme is used having a substantial reverse transcriptase activity at said elevated temperature, said first enzyme being substantially inactive; a composition for use in reverse transcription of RNA, the composition comprising a) a first enzyme having reverse transcriptase activity at temperatures not exceeding 55°C b) a second enzyme having reverse transcriptase activity at elevated temperatures in the range of 45°C - 95°C, said first enzyme has substantially higher activity than said second enzyme in catalyzing reverse transcription at said temperature not exceeding 55°C; use of thermostable enzyme having reverse transcriptase activity in the preparation of a composition for use in reverse transcription of RNA which has previously been in vitro reverse transcribed by another enzyme having reverse transcriptase activity;

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/DK 98/00186

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9318176 A	16-09-1993	US 5262311 A	16-11-1993
		CA 2102784 A	12-09-1993
		EP 0592626 A	20-04-1994
		JP 7500735 T	26-01-1995
		US 5599672 A	04-02-1997
		US 5665547 A	09-09-1997
EP 0743367 A	20-11-1996	DE 19518505 A	21-11-1996
		JP 8308598 A	26-11-1996
WO 9705286 A	13-02-1997	US 5712126 A	27-01-1998
		AU 692403 B	04-06-1998
		AU 6763696 A	26-02-1997
		CA 2201468 A	13-02-1997
		EP 0797685 A	01-10-1997
		JP 10509329 T	14-09-1998
WO 9513369 A	18-05-1995	US 5459037 A	17-10-1995
		AU 687127 B	19-02-1998
		AU 1055195 A	29-05-1995
		AU 6711098 A	09-07-1998
		CA 2174966 A	18-05-1995
		EP 0726946 A	21-08-1996
		FI 962000 A	10-05-1996
		JP 9509306 T	22-09-1997
		NO 961902 A	12-07-1996
		US 5807680 A	15-09-1998
WO 9508647 A	30-03-1995	US 5482845 A	09-01-1996
		AU 7842594 A	10-04-1995
		US 5637685 A	10-06-1997
WO 9729211 A	14-08-1997	AU 2264197 A	28-08-1997